

FUNGICIDE USE IN ALFALFA: WHAT FOUR YEARS OF RESEARCH HAS TAUGHT US

Damon L. Smith ^{1/}, Scott Chapman ^{2/}, Bryan Jensen ^{3/}, Greg Blonde ^{4/}, and Bill Halfman ^{4/}

Introduction

Over the past several years, interest in using foliar-applied fungicides on alfalfa for dairy production has increased. This has subsequently led to new labeling for foliar fungicide products for use on alfalfa. Research at the University of Wisconsin-Madison began in 2011 to evaluate some of the products labeled for use in alfalfa. From 2011 to 2014 replicated on-farm and research station trials were conducted to evaluate the utility of using fungicide on alfalfa for dairy production.

Methods

Trials were located at various locations in each of the research years and included plots in Monroe County, Waupaca County, and Columbia County. Treatments in all trials were replicated four to six times. Each individual plot comprised a minimum area of 400 sq. ft. Treatments were applied using a backpack small-plot sprayer calibrated to deliver 20 gallons of water per acre. All treatments were applied at six to eight inches of growth after each cutting. Applications were made for three cuttings per season. Alfalfa was harvested from each plot for each cutting using a small plot harvester. For some trials foliar disease data were collected. For all trials, quality was evaluated by the University of Wisconsin Soil and Forage Testing Laboratory located in Marshfield Wisconsin. Yield, quality, and disease (where applicable) data were evaluated for each cutting, at each location, for each year.

Results

In total, 35 separate trials (cutting x site x year) were conducted over the four-year period. In the majority of the trials disease levels were low and no significant difference in foliar disease and defoliation was identified between treatments. Some detectable differences in quality were identified between treatments in some trials. However, relative forage quality was typically greater than 150 (Prime Grade) for both treated and non-treated alfalfa. Yield was significantly greater ($\alpha=0.05$) in fungicide treated plots for only 12 of the 35 trials. No particular cutting-timing resulted in a consistent increase in yield when the treatment effect was significant. The average dry matter yield increase over the non-treated control plots was 0.22 tons/acre in trials where fungicide treatment resulted in an increase in yield. The average approximate cost to apply one

^{1/} Assistant Professor, Department of Plant Pathology, 1630 Linden Drive, University of Wisconsin-Madison, Madison, WI, 53706

^{2/} Researcher, Departments of Plant Pathology and Entomology, 1630 Linden Drive, University of Wisconsin-Madison, Madison, WI, 53706

^{3/} Outreach Program Manager, Integrated Pest Management Program, 1630 Linden Drive, University of Wisconsin-Madison, Madison, WI, 53706

^{4/} Agricultural Agent, UW-Extension

fungicide application (fungicide plus custom applicator cost) is estimated to be \$28.00 USD. Considering this cost, the added value per acre for the 12 trials where fungicide increased yield was estimated to be \$13.80 (\$0.10 per pound dry matter hay).

In some cases fungicide can increase alfalfa yield. This is estimated to occur only 34% of the time when fungicide is used. Alfalfa growers are encouraged to focus on timely alfalfa harvest rather than rely on fungicide application to obtain high-quality, high-yielding alfalfa forage.